| | Food A | llergy Action Plan | | |
|---------------------------------|---------------------------------|-----------------------------------|--|-----------------------------------|
| Student's Name: | | D.O.B:Teach | er: | Place |
| ALLERGY | TO: | | | Child's Picture |
| Asthmatic Y | Yes* No * | Higher risk for severe reaction | | Here |
| | ♦ STF | P 1: TREATMENT ◆ | | , |
| Symptoms: | | | Give Checked Medication**: (To be determined by physician authorizing treatment) | |
| If a food a | allergen has been ingested, but | no symptoms: | ☐ Epinephrine | ☐ Antihistamine |
| Mouth | Itching, tingling, or swellin | g of lips, tongue, mouth | ☐ Epinephrine | ☐ Antihistamine |
| Skin | Hives, itchy rash, swelling | of the face or extremities | ☐ Epinephrine | ☐ Antihistamine |
| Gut | Nausea, abdominal cramps | , vomiting, diarrhea | ☐ Epinephrine | ☐ Antihistamine |
| ■ Throat† | Tightening of throat, hoars | eness, hacking cough | ☐ Epinephrine | ☐ Antihistamine |
| ■ Lung† | Shortness of breath, repetit | ive coughing, wheezing | ☐ Epinephrine | ☐ Antihistamine |
| Heart† | Thready pulse, low blood p | ressure, fainting, pale, blueness | ☐ Epinephrine | ☐ Antihistamine |
| Other† | | | ☐ Epinephrine | ☐ Antihistamine |
| If reaction | is progressing (several of the | above areas affected), give | ☐ Epinephrine | ☐ Antihistamine |
| Antihistamiı | ne: give | | | |
| Othom sive | | medication/dose/route | | |
| Other: give_ | | medication/dose/route | | |
| | ◆ <u>S</u> | ΓΕΡ 2: EMERGENCY CAL | <u>LS</u> ♦ | |
| 1. Call 911 (c may be nee | or Rescue Squad: |) . State that an a | allergic reaction has been | treated, and additional epinephri |
| 2. Dr | | at | | |
| 3. Emergency Name/Relation | | Phone Number(s) | | |
| a | | 1.) | 2.) | |
| b | | 1.) | 2.) | |
| c | | 1.) | 2.) | |
| EVEN IF PARE | ENT/GUARDIAN CANNOT BE R | EACHED, DO NOT HESITATE TO M | EDICATE OR TAKE CHII | D TO MEDICAL FACILITY! |
| Parent/Guardia | an Signature | | Date | |
| Doctor's Signa | | | Date | |
| · · | (Required) | | | |

| TRAINED STAFF MEMBERS | | | | |
|-----------------------|------|--|--|--|
| 1 | Room | | | |
| 2 | Room | | | |
| 3 | Room | | | |

EpiPen® and EpiPen® Jr. Directions

Pull off gray activation cap.



 Hold black tip near outer thigh (always apply to thigh).



Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. Remove the EpiPen® unit and massage the injection area for 10 seconds. Twinject™ 0.3 mg and Twinject™ 0.15 mg Directions



- Pull off green end cap, then red end cap.
- Put gray cap against outer thigh, press down firmly until needle penetrates. Hold for 10 seconds, then remove.



SECOND DOSE ADMINISTRATION:

If symptoms don't improve after 10 minutes, administer second dose:

- Unscrew gray cap and pull syringe from barrel by holding blue collar at needle base.
- Slide yellow or orange collar off plunger.
- Put needle into thigh through skin, push plunger down all the way, and remove.



Once EpiPen® or Twinject™ is used, call the Rescue Squad. Take the used unit with you to the Emergency Room. Plan to stay for observation at the Emergency Room for at least 4 hours.

For children with multiple food allergies, consider providing separate Action Plans for different foods.



**Medication checklist adapted from the Authorization of Emergency Treatment form developed by the Mount Sinai School of Medicine. Used with permission.